

# MATH 223: Multivariable Calculus

## Notes on Class 1

September 11, 2023

## CALCULUS: Limits, Derivatives, Integrals of Functions

Classic Setting:  $y = f(x)$  where  $f : \mathcal{R}^1 \rightarrow \mathcal{R}^1$

Input and Output Are Each Single Numbers

Graph is a CURVE (1 Dimensional ) in Plane ( $\mathcal{R}^2$ )

Idea of a function Generalizes Easily:

Input: One Object

Output: One Object

**VECTOR is unifying concept of Linear Algebra and  
Multivariable Calculus**

Calculus I and II: Real-Valued Function of Real Variable

Multivariable Calculus: Vector-Valued Function of Vector Variable

Ultimate Goal:  $f : \mathcal{R}^n \rightarrow \mathcal{R}^m$ .

*Example:* Let  $A$  be a  $3 \times 4$  matrix and  $\mathbf{x}$  a  $4 \times 1$  vector.

Consider the function  $f(\mathbf{x}) = A\mathbf{x}$

$$\begin{pmatrix} - & - & - & - \\ - & - & - & - \\ - & - & - & - \end{pmatrix} \begin{pmatrix} - \\ - \\ - \\ - \end{pmatrix} = \begin{pmatrix} - \\ - \\ - \end{pmatrix}$$

$3 \times 4$                        $4 \times 1$                $3 \times 1$

$A$                                        $\mathbf{x}$                        $A\mathbf{x}$

## Classic Applications of Calculus

- ▶ Motion of Object Along Straight Line  
(Position, Velocity, Acceleration )
- ▶ Profit as a Function of Price
- ▶ Amount of Drug in Bloodstream at time  $t$

## Real World Is Much More Complicated

- ▶ Motion: Objects Move in Plane or Space  
Need Vector to Describe Location
- ▶ Profit: Depends on Prices, Demand, Taxes, Production Costs
- ▶ GPA: Function of Many Course Grades

INPUT: Covid Budget

OUTPUT: Amount for Masks, Vaccine, Hospital Equipment, etc.

## Definition of Derivative

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \text{ IF THE LIMIT EXISTS}$$

*Example:* Find  $f'(x)$  if  $f(x) = x^3$

*Solution:*

$$\begin{aligned} f(x+h) - f(x) &= (x+h)^3 - x^3 \\ &= x^3 + 3x^2h + 3xh^2 + h^3 - x^3 \\ &= 3x^2h + 3xh^2 + h^3 \\ &= h [3x^2 + 3xh + h^2] \end{aligned}$$

$$\text{so } \frac{f(x+h) - f(x)}{h} = [3x^2 + 3xh + h^2] \text{ if } h \neq 0$$

$$\text{Hence } f'(x) = \lim_{h \rightarrow 0} 3x^2 + 3xh + h^2 = 3x^2$$

*Example:* Determine  $g'(x)$  if  $g(x) = x f(x)$  where  $f$  is a differentiable function.

*Solution:*

$$\begin{aligned}g(x+h) - g(x) &= (x+h)f(x+h) - xf(x) \\ &= xf(x+h) + hf(x+h) - xf(x) \\ &= x[f(x+h) - f(x)] + hf(x+h)\end{aligned}$$

Thus The Difference Quotient is

$$\frac{g(x+h) - g(x)}{h} = x \frac{f(x+h) - f(x)}{h} + f(x+h)$$

Taking Limits as  $h \rightarrow 0$ :

$$\frac{f(x+h) - f(x)}{h} \rightarrow f'(x)$$

$f(x+h) \rightarrow f(x)$  since  $f$  is continuous

$$\text{Hence } g'(x) = xf'(x) + f(x)$$