

MATH 223: Multivariable Calculus



Class 23: November 3, 2023

DON'T FORGET TO

FALL
BACK



The Department of Mathematics and Statistics invites you to

Get the Scoop!



...at the **Math/Stat Ice Cream Social**

When: Tuesday, November 7th at 3:45pm

Where: Davis Family Library 105A/B

What: Come and “get the scoop” on spring course offerings within our department while you enjoy some ice cream. Ask faculty members questions about their spring courses and/or get advice on how to build a course of study that suits your needs/interests (while you build an ice cream sundae). Are you a seasoned math/stat major? Please join us and share your advice/experience with other students!

Who should attend: Anyone interested in taking a mathematics or statistics class in Spring 2024, new/current mathematics/statistics majors, and students interested in becoming mathematics/statistics majors!

See you there!



Multiple Integrals
Notes on Assignment 21
Assignment 22

Activity	Date	Approximate Weight
Exam 3	November 29	20%
Project	December 11	10%
Final Exam	December 13 and 14	30%

The Week Ahead:

Iterated Integral (Last Time)
Definition of Multiple Integrals
Properties of the Integral
Change of Variable

Instances of the Integral: I

The Classic Case

$$f := \mathbb{R}^1 \rightarrow \mathbb{R}^1$$

Definite Integral $\int_a^b f(x) dx$

Indefinite Integral $\int f(x) dx$

Instances of the Integral: II

Vector-Valued Functions of a Real Variable

$$f := \mathbb{R}^1 \rightarrow \mathbb{R}^n$$

$$f(t) = [f_1(t), f_2(t), \dots, f_n(t)]$$

$$\int f(t) = \left[\int f_1(t), \int f_2(t), \dots, \int f_n(t) \right]$$


Instances of the Integral: III

Line Integral = Path Integral
(Will Study in Chapter 7)

γ is graph of $g : \mathbb{R}^1 \rightarrow \mathbb{R}^n, a \leq t \leq b$

Force Field $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$

$$\int_{\gamma} F = \int_a^b F(g(t)) \cdot g'(t) dt$$





Instances of the Integral: IV

Iterated Integral

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^1$$

$$\int_a^b \left(\int_{y=g_1(x)}^{y=g_2(x)} f(x, y) dy \right) dx$$



Instances of the Integral: V


The Multiple Integral

A Generalization of Situation 1

$$\int_a^b f(x)dx = \lim_{\max(\Delta(x_i)) \rightarrow 0} \sum_{i=1}^n f(x_i)\Delta x_i$$

Δx_i = length of i th subdivision

x_i = any point in i th subinterval



Instances of the Integral: V


Unequal Divisions

$$\int_a^b f(x) dx = \lim_{\max(\Delta(x_i)) \rightarrow 0} \sum_{i=1}^n f(x_i) \Delta x_i$$


Δx_i = length of i th subdivision

x_i = any point in i th subinterval

Riemann Sums of Unequal Length Subintervals



First Extension: $f : \mathbb{R}^2 \rightarrow \mathbb{R}^1$ on a rectangle \mathcal{R}



Cover \mathcal{R} with a grid G of horizontal and vertical lines
 $mesh(G) = m(G) =$ maximum length of edge of interval in grid.

Number Rectangles R_1, R_2, \dots, R_k with area of Rectangle R_i
denoted by $A(R_i)$.

Pick a point \vec{x}_i in R_i .


$$\text{Form } \sum_{i=1}^k f(\vec{x}_i)A(R_i); \text{ Take } \lim_{m(G) \rightarrow 0} \sum_{i=1}^k f(\vec{x}_i)A(R_i)$$

The limit is the integral of f over \mathcal{R} and is denoted $\int_{\mathcal{R}} f dA$

Second Extension: $f : \mathbb{R}^2 \rightarrow \mathbb{R}^1$ on a BOUNDED set \mathcal{B}

Cover \mathcal{B} with a grid G of horizontal and vertical lines

Let R_1, R_2, \dots, R_k be all bounded rectangles formed by G that lie **inside** \mathcal{B} .



Choose \vec{x}_i in R_i .

$$\text{Take } \lim_{m(G) \rightarrow 0} \sum_{i=1}^k f(\vec{x}_i) A(R_i)$$

The limit is the integral of f over \mathcal{B} and is denoted $\int_{\mathcal{B}} f dV$

Theorem: If $\int_B f dV$ exists and iterated integrals exist for some orders of partial integration, then all of these integrals are equal.

Proof:

Apostol, *Mathematical Analysis*

Sprivak, *Calculus on Manifolds*



When Does The Integral Exist?


Idea: f does not have too many points of discontinuity

Definition: A set S has **zero content** if $\int_S 1dV = 0$.

Theorem: Let \mathcal{B} be a bounded set in \mathbb{R}^n whose boundary has zero content. Let f be a bounded function bounded on \mathcal{B} . If f is continuous on \mathcal{B} except perhaps on a set of zero content, then $\int_{\mathcal{B}} fdV$ exists.

Example from Calculus 1

$$f(x) = \begin{cases} 1 & 0 \leq x \leq 1, \\ 2 & 1 < x \leq 2 \end{cases}$$



$$\int_1^2 f(x) dx = 3$$

Generalize For a Function $f : \mathbb{R}^n \rightarrow \mathbb{R}^1$

Coordinate Rectangle in \mathbb{R}^n

$$\mathcal{R} = \{(x_1, x_2, \dots, x_n) : a_1 \leq x_1 \leq b_1, a_2 \leq x_2 \leq b_2, \dots, a_n \leq x_n \leq b_n\}$$

Volume or Content of \mathcal{R}

$$V(\mathcal{R}) = (b_1 - a_1)(b_2 - a_2) \dots (b_n - a_n)$$

Grid: a finite set of $n - 1$ =dimensional planes in \mathbb{R} parallel to the coordinate planes.

G divides \mathbb{R} into a finite number of bounded "rectangles"

$\mathcal{R}_1, \mathcal{R}_2, \dots, \mathcal{R}_k$ and possibly other unbounded "rectangles,"

The mesh $m(G)$ of a grid = maximum "length" of a side of the rectangles $\mathcal{R}_1, \mathcal{R}_2, \dots, \mathcal{R}_k$

A set \mathcal{B} is bounded if it can be covered by a grid.

$$\text{Then } \int_{\mathcal{B}} f dV = \lim_{m(G) \rightarrow 0} \sum_{i=1}^k f(\vec{x}_i) V(R_i)$$

if this limit exists. for all grids and all choices of \vec{x}_i in R_i .

$$\text{Then } \int_{\mathcal{B}} f dV = \lim_{m(G) \rightarrow 0} \sum_{i=1}^k f(\vec{x}_i) V(R_i)$$

if this limit exists. for all grids and all choices of \vec{x}_i in R_i .

$$\text{Content of } \mathcal{B} = \int_{\mathcal{B}} 1 dV = \begin{cases} \text{Length of } \mathcal{B} & \text{if } \mathcal{B} \subset R^1, \\ \text{Area of } \mathcal{B} & \text{if } \mathcal{B} \subset R^2 \\ \text{Volume of } \mathcal{B} & \text{if } \mathcal{B} \subset R^3 \end{cases}$$

Example Evaluate $\int_{\mathcal{B}}(x^2 + 5y)dV$ where $0 \leq x \leq 1, 0 \leq y \leq 3$
using the definition.

The existence of the integral is guaranteed since \mathcal{B} is bounded and
 $f(x, y) = x^2 + 5y$ is continuous on \mathcal{B}

Hence any sequence of Riemann sums with mesh going to 0 can be
used.

For each $n = 1, 2, \dots$ consider the Grid G_n consisting of
the vertical lines $x = \frac{i}{n}, i = 0, 1, \dots, n$ and
the horizontal lines $y = \frac{j}{n}, j = 0, 1, \dots, 3n$

Then mesh of $G_n = \frac{1}{n}$ and Area of Rectangle $R_{ij} = \frac{1}{n^2}$

$$\text{Riemann sum is } \sum_{i=1}^n \left(\sum_{j=1}^{3n} \left[\left(\frac{i}{n} \right)^2 + 5 \left(\frac{j}{n} \right) \right] \right) A(R_{ij})$$

$$= \frac{1}{n^2} \left[\sum_{i=1}^n \sum_{j=1}^{3n} \left(\frac{i}{n} \right)^2 + \sum_{i=1}^n \sum_{j=1}^{3n} 5 \left(\frac{j}{n} \right) \right]$$

$$= \frac{1}{n^2} \left[3n \sum_{i=1}^n \left(\frac{i}{n} \right)^2 + n \sum_{j=1}^{3n} \frac{5j}{n} \right]$$

$$= \frac{1}{n^2} \left[\frac{3n}{n^2} \sum_{i=1}^n i^2 + \frac{5n}{n} \sum_{j=1}^{3n} j \right]$$


$$= \frac{1}{n^2} \left[\frac{3}{n} \frac{n(n+1)(2n+1)}{6} + 5 \frac{(3n)(3n+1)}{2} \right]$$

$$\text{Riemann sum is } \sum_{i=1}^n \left(\sum_{j=1}^{3n} \left[\left(\frac{i}{n}\right)^2 + 5\left(\frac{j}{n}\right) \right] \right) A(R_{ij})$$

$$\begin{aligned} &= \frac{1}{n^2} \left[\frac{1}{2}(n+1)(2n+1) + \frac{15}{2}n(3n+1) \right] \\ &= \frac{1}{2} \left[\left(1 + \frac{1}{n}\right)\left(2 + \frac{1}{n}\right) \right] + \frac{15}{2} \left[3 + \frac{1}{n} \right] \end{aligned}$$

$$\text{Hence } \lim_{n \rightarrow \infty} = \frac{1}{2}(2) + \frac{15}{2}(3) = \frac{47}{2}$$

**Pooh?
Yeah Piglet?
I'm tired of all this.
I am too Piglet. I am too.**



**There Must Be a Better
Way!**

Evaluate As Iterated Integral

$$\int_{x=0}^{x=1} \int_{y=0}^{y=3} (x^2 + 5y) dy dx$$

$$= \int_{x=0}^{x=1} \left[x^2 y + \frac{5}{3} y^2 \right]_{y=0}^{y=3} dx$$

$$= \int_0^1 3x^2 + \frac{45}{2} dx = \left[x^3 + \frac{45}{2} x \right]_0^1 = \left(1 + \frac{45}{2} \right) - (0 + 0) = \frac{47}{2}$$