


MATH 223: Multivariable Calculus



Class 25: November 8, 2023



Notes on Assignment 23
Assignment 24
Jacobi's Theorem on Change of Variable

Announcements

Review Improper Integrals:

$$\int_1^{\infty} \frac{1}{x^n} dx$$

Progress Report on Location Problem:

Due By Friday, November 17

Should Have Explicit Function To Minimize With Full Rationale

This Week:
Change of Variable
Leibniz Rule
Improper Integrals
Application to Probability

Leibniz Rule: Interchanging Differentiation and Integration

If g_y is continuous on $a \leq x \leq b, c \leq y \leq d$, then

$$\frac{d}{dy} \int_a^b g(x, y) dx = \int_a^b \frac{\partial}{\partial y} g(x, y) dx$$

$$\frac{d}{dy} \int_a^b g(x, y) dx = \int_a^b \frac{\partial}{\partial y} g(x, y) dx$$

Example Compute $f(x) = \int_0^1 \frac{u^x - 1}{\ln u} du$

By Leibniz:

$$f'(x) = \int_0^1 \frac{1}{\ln u} (u^x \ln u) du = \int_0^1 u^x du = \frac{u^{x+1}}{x+1} \Big|_{u=0}^{u=1} = \frac{1}{x+1}$$

So $f(x) = \ln(x+1) + C$ for some constant C .

To Find C , evaluate at $x = 0$:

$$f(0) = \int_0^1 \frac{u^0 - 1}{\ln u} du = \int_0^1 0 = 0$$

But $f(0) = \ln(0+1) + C = \ln(1) + C = 0 + C = C$ so $C = 0$ and

$$f(x) = \ln(x+1)$$

Example: Find $f'(y)$ if $f(y) = \int_0^1 (y^2 + t^2) dt$

Method I: $f(y) = \int_0^1 (y^2 + t^2) dt = (y^2 t + \frac{t^3}{3}) \Big|_{t=0}^{t=1} = y^2 + \frac{1}{3}$ so
 $f'(y) = 2y$

Method II: (Leibniz) $f'(y) = \int_0^1 2y dt = 2yt \Big|_0^1 = 2y$

Proof of Leibniz Rule

To Show:

$$\frac{d}{dy} \int_a^b g(x, y) dx = \int_a^b \frac{\partial}{\partial y} g(x, y) dx$$

Let $f(y) = \int_a^b g(x, y) dx$ and Use Definition of Derivative

$$f'(y) = \lim_{h \rightarrow 0} \frac{f(y+h) - f(y)}{h}$$

$$\frac{f(y+h) - f(y)}{h} = \frac{\int_a^b g(x, y+h) dx - \int_a^b g(x, y) dx}{h} = \frac{\int_a^b (g(x, y+h) - g(x, y)) dx}{h}$$

$$f'(y) = \lim_{h \rightarrow 0} \frac{f(y+h) - f(y)}{h} = \lim_{h \rightarrow 0} \frac{\int_a^b [g(x, y+h) - g(x, y)] dx}{h}$$

Interchange Limit and Integral:

$$= \int_a^b \left(\lim_{h \rightarrow 0} \frac{[g(x, y+h) - g(x, y)]}{h} \right) dx$$

$$= \int_a^b \frac{\partial g}{\partial y}(x, y) dx$$

Alternate Proof of Leibniz Rule

(Uses Iterated Integral)

Begin with $\int_a^b g_y(x, y) dx$

Let $I = \int_c^y (\int_a^b g_y(x, y) dx) dy$

Switch Order of Integration: $I = \int_a^b (\int_c^y g_y(x, y) dy) dx$

$$\begin{aligned} I &= \int_a^b g(x, y) \Big|_{y=c}^{y=y} dx = \int_a^b g(x, y) - g(x, c) dx \\ &= \int_a^b g(x, y) dx - \int_a^b g(x, c) dx \end{aligned}$$

The left term is a function of y and the second is a constant C

Alternate Proof of Leibniz Rule (Continued)

$$I = \int_c^y \left(\int_a^b g_y(x, y) dx \right) dy = \int_a^b g(x, y) dx - C$$


Now Take the Derivative of Each Side with Respect to y , using the Fundamental Theorem of Calculus on the left:

$$\int_a^b g_y(x, y) dx = \frac{d}{dy} \int_a^b g(x, y) dx - 0$$

Richard Feynman

May 11, 1918 – February 15, 1988

Nobel Prize in Physics, 1965



"I used that one damn tool again and again."

" I caught on how to use that method, and I used that one damn tool again and again. [If] guys at MIT or Princeton had trouble doing a certain integral, [then] I come along and try differentiating under the integral sign, and often it worked. So I got a great reputation for doing integrals, only because my box of tools was different from everybody else's, and they had tried all their tools on it before giving the problem to me. (*Surely You're Joking, Mr. Feynman!*)

Richard Feynman's Integral Trick

Change of Variable aka Method of Substitution

A common technique in the evaluation of integrals is to make a change of variable in the hopes of simplifying the problem of determining an antiderivatives



Example: Evaluate $\int_{x=0}^{x=2} \frac{2x}{1+x^2} dx$

$$\begin{array}{l} \text{Let } u = 1 + x^2 \quad \left| \quad x = 0 \rightarrow u = 1 + 0^2 = 1 \right. \\ \text{The } du = 2x dx \quad \left| \quad x = 2 \rightarrow u = 1 + 2^2 = 5 \right. \end{array}$$

$$\int_{x=0}^{x=2} \frac{2x}{1+x^2} dx = \int_{u=1}^{u=5} \frac{1}{u} du = \ln 5 - \ln 1 = \ln 5$$

$$\int_{x=0}^{x=2} \frac{2x}{1+x^2} dx = \int_{u=1}^{u=5} \frac{1}{u} du$$

Let's look at what is happening geometrically:





Not only does the function change, but also the region of integration.

The region of integration changes from an interval of length 2 to an interval of length 4.

The interval also moves to a new location.

In computing multiple integrals, the corresponding change in the region may be more complicated.

By a **change of variable**, we will mean a vector function T from \mathbb{R}^n to \mathbb{R}^n . It is convenient to use different letters to denote the spaces; e.g, $T : U^n \rightarrow \mathbb{R}^n$



Carl Gustav Jacob Jacobi

December 10, 1804 – February 18, 1851



Mathematics exists solely for
the honour of the human mind.

~ Carl Gustav Jacob Jacobi

AZ QUOTES

For further information see his [Biography](#)

Jacobi's Theorem

Let \mathcal{R} be a set in \mathbb{U}^n and $T(\mathcal{R})$ its image under T ; that is,

$$T(\mathcal{R}) = \{T(\vec{u}) : \vec{u} \text{ is in } \mathcal{R}\}$$

Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^1$ is a real-valued function.

Then, under suitable conditions,

$$\int_{T(\mathcal{R})} f(\vec{x}) dV_{\vec{x}} = \int_{\mathcal{R}} f(T(\vec{u})) |\det T'(\vec{u})| dV_{\vec{u}}$$

- ▶ T is continuous differentiable
- ▶ Boundary of \mathcal{R} is finitely many smooth curves
- ▶ T is one-to-one on interior of \mathcal{R}
- ▶ The Jacobian Determinant $\det T'$ is non zero on interior of \mathcal{R} .
- ▶ The function f is bounded and continuous on $T(\mathcal{R})$

$$\int_{T(\mathcal{R})} f(\vec{x}) dV_{\vec{x}} = \int_{\mathcal{R}} f(T(\vec{u}) | \det T'(\vec{u}) |) dV_{\vec{u}}$$

In our example: $u = 1 + x^2$ so $x = \sqrt{u-1}$

Thus $T(u) = \sqrt{u-1} = (u-1)^{1/2}$ so

$$T'(u) = \frac{1}{2}(u-1)^{-1/2} = \frac{1}{2\sqrt{u-1}}$$


$$\int_0^2 \frac{2x}{1+x^2} dx = \int_{T(\mathcal{R})} f(\vec{x}) dV_{\vec{x}} = \int_1^5 f(T(u) | \det T'(u) |) du$$

$$\text{Now } f(T(\vec{u})) = \frac{2T(u)}{1+(T(u))^2} = \frac{2\sqrt{u-1}}{1+u-1} = \frac{2\sqrt{u-1}}{u}$$

$$\det T'(u) = \left| \frac{1}{2\sqrt{u-1}} \right| = \frac{1}{2\sqrt{u-1}} \text{ so } f(T(\vec{u})) \det T'(u) = \frac{1}{u}$$

$$\text{so } \int_0^2 \frac{2x}{1+x^2} dx = \int_1^5 \frac{2\sqrt{u-1}}{u} \frac{1}{2\sqrt{u-1}} du = \int_1^5 \frac{1}{u} du$$

Example: Polar Coordinate Change of Variable

$$\mathcal{U}^2 \quad T \rightarrow \quad \mathcal{R}^2$$


$x = r \cos \theta$
 $y = r \sin \theta$

$$T(r, \theta) = (r \cos \theta, r \sin \theta)$$


$$T' = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} \text{ so } \det T' = r \cos^2 \theta + r \sin^2 \theta = r$$

$$\text{Thus } \int_{T(R)} f(x, y) dx dy = \int_R f(r \cos \theta, r \sin \theta) r dr d\theta$$

$$\int_{T(R)} f(x, y) dx dy = \int_R f(r \cos \theta, r \sin \theta) r dr d\theta$$

Example: $f(x, y) = x^2 + y^2$


$$T(R) = \text{Half Disk} = \{(x, y) : -1 \leq x \leq 1, 0 \leq y \leq \sqrt{1-x^2}\}$$





$$I = \int_{-1}^1 \int_0^{\sqrt{1-x^2}} (x^2 + y^2) dy dx$$

Describe Region in Polar Coordinates: $0 \leq r \leq 1, 0 \leq \theta \leq \pi$

$$I = \int_{\theta=0}^{\pi} \int_{r=0}^1 r^2 r dr d\theta = \int_{\theta=0}^{\pi} \left. \frac{r^4}{4} \right|_0^1 d\theta = \int_{\theta=0}^{\pi} \frac{1}{4} d\theta = \frac{\pi}{4}$$



Look At This Transformation More Closely



$$\begin{aligned} A : 0 \leq r \leq 1, \theta = 0 \\ x = r \cos \theta = r \cos 0 = r \\ y = r \sin \theta = r \sin 0 = 0 \end{aligned}$$

$$\begin{aligned} B : r = 1, 0 \leq \theta \leq \pi \\ x = r \cos \theta = \cos \theta \\ y = r \sin \theta = \sin \theta \end{aligned}$$


$$\begin{aligned} C : 0 \leq r \leq 1, \theta = \pi \\ x = r \cos \theta = r \cos \pi = -r \\ y = r \sin \theta = r \sin \pi = 0 \end{aligned}$$

$$\begin{aligned} D : r = 0, 0 \leq \theta \leq \pi \\ x = r \cos \theta = 0 \\ y = r \sin \theta = 0 \end{aligned}$$

Problem: Evaluate $\iiint_C \sqrt{x^2 + y^2 + z^2} dV$

where C is the ice cream cone

$$\{(x, y, z) : x^2 + y^2 + z^2 \leq 1, x^2 + y^2 \leq \frac{z^2}{3}, z \geq 0\}$$



Example: Spherical Coordinates

$$x = r \sin \phi \cos \theta \quad T : (r, \phi, \theta) \rightarrow (x, y, z)$$

$$y = r \sin \phi \sin \theta \quad \det T' = r^2 \sin \phi$$

$$z = r \cos \phi$$

Problem: Evaluate $\iiint_C \sqrt{x^2 + y^2 + z^2} dV$

where C is the ice cream cone

$$\{(x, y, z) : x^2 + y^2 + z^2 \leq 1, x^2 + y^2 \leq \frac{z^2}{3}, z \geq 0\}$$

$$z \geq 0 \text{ implies } \phi \leq \frac{\pi}{2}$$


$$x^2 + y^2 + z^2 \leq 1 \text{ implies } r \leq 1$$

$$x^2 + y^2 \leq \frac{z^2}{3} \text{ implies } r^2 \sin^2 \phi \leq \frac{r^2 \cos^2 \phi}{3}$$

$$\text{implies } \tan^2 \phi \leq \frac{1}{3} \text{ implies } \phi \leq \frac{\pi}{6}$$

$$\int_{\theta=0}^{2\pi} \int_{\phi=0}^{\pi/6} \int_{r=0}^1 \sqrt{r^2} r^2 \sin \phi \, dr \, d\phi \, d\theta$$

Example: Evaluate $\iiint_D z^2 dV$ where D is the interior of the ellipsoid $\frac{x^2}{4} + \frac{y^2}{16} + \frac{z^2}{9} = 1$



STEP 1: Let $u = \frac{x}{2}$, $v = \frac{y}{4}$, $w = \frac{z}{3}$.

Equation of the ellipsoid becomes $u^2 + v^2 + w^2 = 1$ (unit sphere)

So $x = 2u$, $y = 4v$, $z = 3w$ gives $T(u, v, w) = (2u, 4v, 3w)$ and

$$T' = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix} \text{ so } \det T' = 2 \times 4 \times 3 = 24$$

Thus $\iiint_D z^2 = \iiint (3w)^2 (24) du dv dw = 216 \iiint w^2 du dv dw$

STEP 2: Switch to Spherical Coordinates:

$$u = r \sin \phi \cos \theta, v = r \sin \phi \sin \theta, w = r \cos \phi$$

$$\begin{aligned} 216 \iiint w^2 du dv dw &= 216 \iiint (r \cos \phi)^2 r^2 \sin \phi dr d\phi d\theta \\ &= 216 \int_{\theta=0}^{2\pi} \int_{\phi=0}^{\pi} \int_{r=0}^1 r^4 \cos^2 \phi \sin \phi dr d\phi d\theta \\ &= (216)(2\pi) \int_{\phi=0}^{\pi} \int_{r=0}^1 r^4 \cos^2 \phi \sin \phi dr d\phi \\ &= (216)(2\pi) \frac{1}{5} \int_{\phi=0}^{\pi} \cos^2 \phi \sin \phi d\phi \\ &= \frac{(216)(2\pi)}{5} \left[-\frac{\cos^3 \phi}{3} \right]_{\phi=0}^{\pi} = \frac{(216)(2\pi)}{5} \frac{2}{3} = \frac{288\pi}{5} \end{aligned}$$