

MATH 223: Multivariable Calculus

Notes on Class 3

September 15, 2023

Derivatives and Integrals for $\mathbf{f} : \mathcal{R}^1 \rightarrow \mathcal{R}^n$

Vector-Valued Functions of a Real Variable

Begin with $\mathbf{F} : \mathcal{R}^1 \rightarrow \mathcal{R}^2$

$$\mathbf{F}(x) = (f(x), g(x))$$

Difference Quotient

$$\frac{\mathbf{F}(x+h) - \mathbf{F}(x)}{h} = \left(\frac{f(x+h) - f(x)}{h}, \frac{g(x+h) - g(x)}{h} \right)$$

So

$$\mathbf{F}'(x) = \lim_{h \rightarrow 0} \frac{\mathbf{F}(x+h) - \mathbf{F}(x)}{h} = (f'(x), g'(x))$$

Example: $\mathbf{F}(x) = (\cos x, x^3 - 2x)$

Solution: $\mathbf{F}'(x) = (-\sin x, 3x^2 - 2)$

Example: $\mathbf{F}(t) = (\tan t, \ln t)$

Solution: $\mathbf{F}'(t) = (\sec^2 t, \frac{1}{t})$

Nothing Special about $m = 2$

$$\mathbf{F}(x) = (f_1(x), f_2(x), \dots, f_m(x))$$

$$\mathbf{F}'(x) = (f_1'(x), f_2'(x), \dots, f_m'(x))$$

Example: $\mathbf{F}(t) = (t^7, t^{-3}, \sin(t^2))$

Derivative: $(7t^6, -3t^{-4}, 2t \cos(t^2))$

IMAGE of \mathbf{F} is a Curve (1 dimensional) in \mathcal{R}^m .

Tangent Lines

$$\mathbf{L}(t) = \mathbf{F}(x) + t\mathbf{F}'(x)$$

Example: $\mathbf{F}(x) = (x^3 + 7x + 3, 8 + \sin x)$

Then $\mathbf{F}'(x) = (3x^2 + 7, \cos x)$

At $x = 0$: $\mathbf{F}(0) = (3, 8)$ and $\mathbf{F}'(0) = (7, 1)$

The Equation for the tangent line at $(3,8)$ is

$$\mathbf{L}(t) = (3, 8) + t(7, 1) = (3 + 7t, 8 + t)$$

We can write as $x = 3 + 7t, y = 8 + t$ so

$$t = \frac{x-3}{7} \text{ and } y = 8 + \frac{x-3}{7}$$

Bottom Line

$$\mathbf{F} = (f_1, f_2, \dots, f_m)$$

where each $f_i : \mathcal{R}^1 \rightarrow \mathcal{R}^1$

\mathbf{F} is continuous if and only if each f_i is continuous

\mathbf{F} is differentiable if and only if each f_i is differentiable

$$\mathbf{F}' = (f_1', f_2', \dots, f_m')$$

$$\int \mathbf{F} = \left(\int f_1, \int f_2, \dots, \int f_m \right)$$

KEY STEP IS THEOREM 2.2.1