


MATH 223: Multivariable Calculus



Class 9
Friday, September 29, 2023



- ▶ Notes on Assignment 8
- ▶ Assignment 9

Announcements

Exam 1: Next Wednesday, 7 PM -
No Time Limit

No Books, Computers, Smartphones, etc.

One Page of Notes OK

Focus on Chapters 2 and 3

Tangent Planes To Surfaces

(I) $f : \mathcal{R}^2 \rightarrow \mathcal{R}^1$, **a a point in \mathcal{R}^2**

Tangent plane to graph of f at $(\mathbf{a}, f(\mathbf{a}))$:

$$T(\mathbf{x}) = f(\mathbf{a}) + \nabla f(\mathbf{a}) \cdot (\mathbf{x} - \mathbf{a})$$

(II): $f : \mathcal{R}^2 \rightarrow \mathcal{R}^3$

$$\sigma(s, t) = (f(s, t), g(s, t), h(s, t))$$

$$\sigma_s(s, t) = (f_s, g_s, h_s) \text{ and } \sigma_t(s, t) = (f_t, g_t, h_t)$$

Tangent Plane at $\sigma(\mathbf{a})$:


$$\sigma(\mathbf{a}) + (s, t) \begin{pmatrix} f_s(\mathbf{a}) & g_s(\mathbf{a}) & h_s(\mathbf{a}) \\ f_t(\mathbf{a}) & g_t(\mathbf{a}) & h_t(\mathbf{a}) \end{pmatrix}$$

Note: $1 \times 3 + (1 \times 2)(2 \times 3)$

Writing vectors vertically: $\sigma = \begin{pmatrix} f \\ g \\ h \end{pmatrix}$, $\sigma' = \begin{pmatrix} f' \\ g' \\ h' \end{pmatrix}$

Tangent Plane: $T \begin{pmatrix} s \\ t \end{pmatrix} = \sigma(\mathbf{a}) + \sigma'(\mathbf{a}) \begin{pmatrix} s \\ t \end{pmatrix}$

Parametrized Surfaces




Function from	$\mathcal{R}^2 \rightarrow \mathcal{R}^3$
Domain	Patch in Plane
Image	Surface in Space
Graph	Lives in \mathcal{R}^5

Need for Parametrizations: Graph of $f : \mathcal{R}^1 \rightarrow \mathcal{R}^1$ is a curve but not every curve is the graph of such a function

Similarly, graph of $f : \mathcal{R}^2 \rightarrow \mathcal{R}^1$ is a surface but not every surface is the graph of such a function.

Example: $\sigma(s, t) = (s \cos t, s \sin t, t), 0 \leq s \leq 4, 0 \leq t \leq 3\pi$




Point: $(1, \pi/4)$ so $\sigma(1, \pi/4) = \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, \frac{\pi}{4}\right)$

$\sigma_s(s, t) = (\cos t, \sin t, 0)$ and $\sigma_t(s, t) = (-s \sin t, s \cos t, 1)$


At $(1, \frac{\pi}{4})$, representation of the tangent plane is

$$\sigma\left(1, \frac{\pi}{4}\right) + \sigma_s\left(1, \frac{\pi}{4}\right) s + \sigma_t\left(1, \frac{\pi}{4}\right) t$$



Parametrize Unit Sphere

$$\sigma(s, t) = (\cos t \cos s, \sin t \cos s, \sin s), 0 \leq s \leq 2\pi, 0 \leq t \leq 2\pi$$



$$x = \cos t \cos s, y = \sin t \cos s, z = \sin s$$


$$\begin{aligned}x^2 + y^2 + z^2 &= \cos^2 t \cos^2 s + \sin^2 t \cos^2 s + \sin^2 s \\&= \cos^2 s (\cos^2 t + \sin^2 t) + \sin^2 s \\&= \cos^2 s + \sin^2 s = 1\end{aligned}$$

Utility

Utility = happiness, satisfaction, pleasure, usefulness

$$u(x), x \geq 0$$

Typical Assumptions: u is increasing, concave down function
("decreasing returns to scale")



Example: $u(x) = x^{1/3}$ so $u'(x) = \frac{1}{3x^{2/3}}$, $u''(x) = -\frac{2}{9}x^{-5/3}$

Example: 2 Goods with $u(x, y) = \sqrt[3]{xy}$

Each unit of x costs \$35 and each unit of y costs \$80

We have \$ D to spend: Budget Constraint: $35x + 80y = D$

Goal: Maximize Utility:

$$80y = D - 35x \text{ so } y = \frac{D - 35x}{80}$$

$$u(x, y) = f(x) = \sqrt[3]{\frac{x(D - 35x)}{80}}$$


f is maximized when $\frac{x(D-35x)}{80}$ is maximized.

$G(x) = x(D - 35x) = Dx - 35x^2$. has $G'(x) = D - 70x$ and

$G''(xx) = -70$ Hence there is a maximum when $x = D/70$

$$\text{Then } y = \frac{D - 35(D/70)}{80} = D/160$$

Clairaut's Theorem on Equality of Mixed Partial
If f_{xy} and f_{yx} are continuous at \mathbf{a} , then $f_{xy}(\mathbf{a}) = f_{yx}(\mathbf{a})$



May 7, 1713 – May 17, 1765

Clairaut's Theorem on Equality of Mixed Partial

If f_{xy} and f_{yx} are continuous at \mathbf{a} , then $f_{xy}(\mathbf{a}) = f_{yx}(\mathbf{a})$

$$f(x, y) = \begin{cases} 2xy \frac{x^2 - y^2}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$$

It Turns Out That

$$f_{xy}(0, 0) = -2$$

$$f_{yx}(0, 0) = +2$$

Mixed Partial Are Not Equal