

MATH 223

Some Notes on Assignment 12

Exercises 26, 27, 28, and 29 in Chapter 4. (Old Exercises 19, 20, 21, 22)

26. Find the directional derivative of $f(x, y) = x^3\sqrt{y}$ at (7,25) in the direction toward (3,-4).

Solution: If $f(x, y) = x^3\sqrt{y}$, then the gradient, $\nabla f(x, y) = [3x^2\sqrt{y}, \frac{x^3}{2\sqrt{y}}]$ is the 1 by 2 vector containing the partial derivatives of f with respect to x and y . To find the derivative of f in the direction of $\mathbf{v} = (3, 4)$ we need only find the product $\nabla f(7, 25) \cdot \mathbf{u}$ where \mathbf{u} is the unit vector with the same direction as \mathbf{v} .

$$\mathbf{v} = (3, -4) \rightarrow \mathbf{u} = (3, -4)\left(\frac{1}{\sqrt{3^2 + 4^2}}\right) = \left(\frac{3}{5}, \frac{-4}{5}\right)$$

$$f_{\mathbf{u}}(7, 25) = \nabla f(7, 25)\mathbf{u} = \left[735, \frac{343}{10}\right]\left(\frac{3}{5}, \frac{-4}{5}\right) = \frac{10339}{25} = 413.56$$

27: Find the directional derivative of $f(x, y, z) = xy^2z^3$ at (1,6,2) in the direction toward (3, -1, -1).

Solution: To find a derivative of $f(x, y, z)$ in the direction of $\mathbf{v} = (3, -1, -1)$, we must find the gradient of f at (1, 6, 2) and a unit vector in the direction of \mathbf{v} . If we take the partial derivatives with respect to each variable of f we find the gradient to be $\nabla f(x, y, z) = [y^2z^3, 2xyz^3, 3xy^2z^2]$. Evaluated at the point (1, 6, 2) the gradient is [288, 96, 432]. The unit vector in the direction of v is $\frac{\mathbf{v}}{|\mathbf{v}|} = \left(\frac{3}{\sqrt{11}}, \frac{-1}{\sqrt{11}}, \frac{-1}{\sqrt{11}}\right)$. From Theorem 4.3.1. we have

$$f_{\mathbf{u}}(1, 6, 2) = [288, 96, 432]\left(\frac{3}{\sqrt{11}}, \frac{-1}{\sqrt{11}}, \frac{-1}{\sqrt{11}}\right) = \frac{336}{\sqrt{11}} \approx 101.308$$

28: At the point (3,2), a certain function f has a directional derivative of $\frac{288}{5}$ in the direction (-3,4) and a directional derivative of $-\frac{36}{13}$ in the direction (-12,5). Find the gradient of f at (3,2).

Solution: Let $\mathbf{v}_1 = (-3, 4)$, and $\mathbf{v}_2 = (-12, 5)$. If the directional derivative of $f(x, y)$ at (3, 2) is $\frac{288}{5}$ in the direction of \mathbf{v}_1 , and $-\frac{36}{13}$ in the direction of v_2 then we have

$$\nabla f(3, 2) \cdot \frac{\mathbf{v}_1}{|\mathbf{v}_1|} = \frac{288}{5},$$

$$\nabla f(3, 2) \cdot \frac{\mathbf{v}_2}{|\mathbf{v}_2|} = \frac{-36}{13}.$$

If we calculate the value of the unit vectors $\frac{\mathbf{v}}{|\mathbf{v}|}$, and expand the gradient of $f(3, 2)$ to be the 1 by 2 matrix of partial derivatives we get

$$[f_x(3, 2), f_y(3, 2)]\left(\frac{-3}{5}, \frac{4}{5}\right) = \frac{288}{5},$$

$$[f_x(3, 2), f_y(3, 2)]\left(\frac{-12}{13}, \frac{5}{13}\right) = \frac{-36}{13},$$

$$\begin{aligned} -3f_x(3, 2) + 4f_y(3, 2) &= 288, \\ -12f_x(3, 2) + 5f_y(3, 2) &= -36. \end{aligned}$$

These last two equations can be rewritten as a system of linear equations of the form

$$\begin{pmatrix} -3 & 4 \\ -12 & 4 \end{pmatrix} \begin{pmatrix} f_x(3, 2) \\ f_y(3, 2) \end{pmatrix} = \begin{pmatrix} 288 \\ -36 \end{pmatrix}$$

Using either a linear algebra software or row reduction we find $f_x(3, 2) = 48$ and $f_y(3, 2) = 108$. The gradient of $f(3, 2)$ is then $\nabla f(3, 2) = (48, 108)$.

29 The real-valued function f is differentiable at a certain point \mathbf{x} in \mathbb{R}^3 with the following known directional derivatives: $-1/\sqrt{14}$ in the direction $(-2, 3, 1)$, 0 in direction $(-5, -1, 8)$ and $2\sqrt{3}$ in direction $(1, 1, 1)$. Find the directional derivative in the direction $(3, 4, -5)$.

Solution: If the directional derivative of $f(x, y, z)$ at a particular point is $\frac{-1}{\sqrt{14}}$ in the direction $(-2, 3, 1)$, 0 in the direction $(-5, -1, 8)$, and $2\sqrt{3}$ in the direction $(1, 1, 1)$, by Theorem 4.3.1 and the definition of directional derivatives we have

$$\begin{aligned} \nabla f(\mathbf{x}) \cdot \frac{(-2, 3, 1)}{|(-2, 3, 1)|} &= \frac{-1}{\sqrt{14}}, \\ \nabla f(\mathbf{x}) \cdot \frac{(-5, 1, 8)}{|(-5, 1, 8)|} &= 0, \\ \nabla f(\mathbf{x}) \cdot \frac{(1, 1, 1)}{|(1, 1, 1)|} &= 2\sqrt{3}. \end{aligned}$$

Notice that each of these equations can be simplified by multiplying by the magnitude of the respective direction vectors to get

$$\begin{aligned} \nabla f(\mathbf{x}) \cdot (-2, 3, 1) &= -1, \\ \nabla f(\mathbf{x}) \cdot (-5, 1, 8) &= 0, \\ \nabla f(\mathbf{x}) \cdot (1, 1, 1) &= (2)(\sqrt{3})(\sqrt{3}) = 6 \end{aligned}$$

If the gradient of $f(\mathbf{x})$ is expanded to be the 1 by 3 matrix of partial derivatives at \mathbf{x} the system of equations becomes

$$\begin{aligned} -2f_x(\mathbf{x}) + 3f_y(\mathbf{x}) + f_z(\mathbf{x}) &= -1, \\ -5f_x(\mathbf{x}) - f_y(\mathbf{x}) + 8f_z(\mathbf{x}) &= 0, \\ f_x(\mathbf{x}) + f_y(\mathbf{x}) + f_z(\mathbf{x}) &= 6. \end{aligned}$$

Rewritten in matrix form this system is

$$\begin{pmatrix} -2 & 3 & 1 \\ -5 & -1 & 8 \\ 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} f_x(\mathbf{x}) \\ f_y(\mathbf{x}) \\ f_z(\mathbf{x}) \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ 6 \end{pmatrix}.$$

Using a linear algebra software or row reduction to solve for each partial derivative reveals $f_x(\mathbf{x}) = 3$, $f_y(\mathbf{x}) = 1$, and $f_z(\mathbf{x}) = 1$. The gradient of f at \mathbf{x} is $\nabla f(\mathbf{x}) = (3, 1, 2)$. Then the directional derivative in the direction $(3, 4, -5)$ is $(3, 1, 2) \cdot \frac{(3, 4, -5)}{\sqrt{3^2+4^2+(-5)^2}} = \frac{3}{5\sqrt{2}}$